

Infinite norm decompositions of C^* -algebras

Farkhad Arzikulov

subclass: *Primary* 46L35, 17C65; *Secondary* 47L30

keywords: *C^* -algebra, infinite norm decompositions*

Abstract

In the given article the notion of infinite norm decomposition of a C^* -algebra is investigated. The norm decomposition is some generalization of Peirce decomposition. It is proved that the infinite norm decomposition of any C^* -algebra is a C^* -algebra. C^* -factors with an infinite and a nonzero finite projection and simple purely infinite C^* -algebras are constructed.

INTRODUCTION

In the given article the notion of infinite norm decomposition of C^* -algebra is investigated. It is known that for any projection p of a unital C^* -algebra A the next equality is valid $A = pAp \oplus pA(1-p) \oplus (1-p)Ap \oplus (1-p)A(1-p)$, where \oplus is a direct sum of spaces. The norm decomposition is some generalization of Peirce decomposition. First such infinite decompositions were introduced in [1] by the author.

In this article a unital C^* -algebra A with an infinite orthogonal set $\{p_i\}$ of equivalent projections such that $\sup_i p_i = 1$, and the set $\sum_{ij}^o p_i A p_j = \{a_{ij} : \text{for any indexes } i, j, a_{ij} \in p_i A p_j, \text{ and } \|\sum_{k=1, \dots, i-1} (a_{ki} + a_{ik}) + a_{ii}\| \rightarrow 0 \text{ at } i \rightarrow \infty\}$ are considered. Note that all infinite sets like $\{p_i\}$ are supposed to be countable.

The main results of the given article are the next:

- for any C^* -algebra A with an infinite orthogonal set $\{p_i\}$ of equivalent projections such that $\sup_i p_i = 1$ the set $\sum_{ij}^o p_i A p_j$ is a C^* -algebra with the component-wise algebraic operations, the associative multiplication and the norm.
- there exist a C^* -algebra A and different countable orthogonal sets $\{e_i\}$ and $\{f_i\}$ of equivalent projections in A such that $\sup_i e_i = 1$, $\sup_i f_i = 1$, $\sum_{ij}^o e_i A e_j \neq \sum_{ij}^o f_i A f_j$.
- if A is a W^* -factor of type II_∞ , then there exists a countable orthogonal set $\{p_i\}$ of equivalent projections in A such that $\sum_{ij}^o p_i A p_j$ is a C^* -factor with a nonzero finite and an infinite projection. In this case $\sum_{ij}^o p_i A p_j$ is not a von Neumann algebra.
- if A is a W^* -factor of type III, then for any countable orthogonal set $\{p_i\}$ of equivalent projections in A . The C^* -subalgebra $\sum_{ij}^o p_i A p_j$ is simple and purely infinite. In this case $\sum_{ij}^o p_i A p_j$ is not a von Neumann algebra.
- there exist a C^* -algebra A with an orthogonal set $\{p_i\}$ of equivalent projections such that $\sum_{ij}^o p_i A p_j$ is not an ideal of A .

1. INFINITE NORM DECOMPOSITIONS

Lemma 1. *Let A be a C^* -algebra, $\{p_i\}$ be an infinite orthogonal set of projections with the least upper bound 1 in the algebra A and let $\mathcal{A} = \{\{p_i a p_j\} : a \in A\}$. Then,*

1) *the set \mathcal{A} is a vector space with the next componentwise algebraic operations*

$$\lambda\{p_i a p_j\} = \{p_i \lambda a p_j\}, \lambda \in \mathbb{C}$$

$$\{p_i a p_j\} + \{p_i b p_j\} = \{p_i(a+b)p_j\}, a, b \in A,$$

2) *the algebra A and the vector space \mathcal{A} can be identified in the sense of the next map*

$$\mathcal{I} : a \in A \rightarrow \{p_i a p_j\} \in \mathcal{A}.$$

Proof. Item 1) of the lemma can be easily proved.

Proof of item 2): We assert that \mathcal{I} is a one-to-one map. Indeed, it is clear, that for any $a \in A$ there exists a unique set $\{p_i a p_j\}$, defined by the element a .

Suppose that there exist different elements a and b in A such that $p_i a p_j = p_i b p_j$ for all i, j , i.e. $\mathcal{I}(a) = \mathcal{I}(b)$. Then $p_i(a-b)p_j = 0$ for all i and j . Observe that $p_i((a-b)p_j(a-b)^*) = ((a-b)p_j(a-b)^*)p_i = 0$ and $(a-b)p_j(a-b)^* \geq 0$ for all i, j . Therefore, the element $(a-b)p_j(a-b)^*$ commutes with every projection in $\{p_i\}$.

We prove $(a-b)p_j(a-b)^* = 0$. Indeed, there exists a maximal commutative subalgebra A_o of the algebra A , containing the set $\{p_i\}$ and the element $(a-b)p_j(a-b)^*$. Since $(a-b)p_j(a-b)^*p_i = p_i(a-b)p_j(a-b)^* = 0$ for any i , then the condition $(a-b)p_j(a-b)^* \neq 0$ contradicts the equality $\sup_i p_i = 1$.

Indeed, in this case $p_i \leq 1 - 1/\|(a-b)p_j(a-b)^*\|(a-b)p_j(a-b)^*$ for any i . Since by $(a-b)p_j(a-b)^* \neq 0$ we have $1 > 1 - 1/\|(a-b)p_j(a-b)^*\|(a-b)p_j(a-b)^*$, then we get a contradiction with $\sup_i p_i = 1$. Therefore $(a-b)p_j(a-b)^* = 0$.

Hence, since A is a C^* -algebra, then $\|(a-b)p_j(a-b)^*\| = \|((a-b)p_j)((a-b)p_j)^*\| = \|((a-b)p_j)\| \|(a-b)p_j\| = \|(a-b)p_j\|^2 = 0$ for any j . Therefore $(a-b)p_j = 0$, $p_j(a-b)^* = 0$ for any j . Analogously, we can get $p_j(a-b) = 0$, $(a-b)^*p_j = 0$ for any j . Hence the elements $a-b$, $(a-b)^*$ commute with every projection in $\{p_i\}$. Then there exists a maximal commutative subalgebra A_o of the algebra A , containing the set $\{p_i\}$ and the element $(a-b)(a-b)^*$. Since $p_i(a-b)(a-b)^* = (a-b)(a-b)^*p_i = 0$ for any i , then the condition $(a-b)(a-b)^* \neq 0$ contradicts the equality $\sup_i p_i = 1$.

Therefore, $(a-b)(a-b)^* = 0$, $a-b = 0$, i.e. $a = b$. Thus the map \mathcal{I} is one-to-one. \square

Lemma 2. *Let A be a C^* -algebra, $\{p_i\}$ be an infinite orthogonal set of projections with the least upper bound 1 in the algebra A and $a \in A$. Then, if $p_i a p_j = 0$ for all i, j , then $a = 0$.*

Proof. Let $p \in \{p_i\}$. Observe that $p_i a p_j a^* = p_i(a p_j a^*) = a p_j a^* p_i = (a p_j a^*)p_i = 0$ for all i, j and $a p_j a^* = a p_j p_j a^* = (a p_j)(p_j a^*) = (a p_j)(a p_j)^* \geq 0$. Therefore, the element $a p_j a^*$ commutes with all projections of the set $\{p_i\}$.

We prove $a p_j a^* = 0$. Indeed, there exists a maximal commutative subalgebra A_o of the algebra A , containing the set $\{p_i\}$ and the element $a p_j a^*$. Since $p_i(a p_j a^*) = (a p_j a^*)p_i = 0$ for any i , then the condition $a p_j a^* \neq 0$ contradicts the equality $\sup_i p_i = 1$ (see the proof of lemma 1). Hence $a p_j a^* = 0$.

Hence, since A is a C^* -algebra, then $\|a p_j a^*\| = \|(a p_j)(a p_j)^*\| = \|(a p_j)\| \|(a p_j)^*\| = \|a p_j\|^2 = 0$ for any j . Therefore $a p_j = 0$, $p_j a^* = 0$ for any j . Analogously we have $p_j a = 0$, $a^* p_j = 0$ for any j . Hence the elements a , a^* commute with all projections

of the set $\{p_i\}$. Then there exists a maximal commutative subalgebra A_o of the algebra A , containing the set $\{p_i\}$ and the element aa^* . Since $p_i aa^* = aa^* p_j = 0$ for any i , then the condition $aa^* \neq 0$ contradicts the equality $\sup_i p_i = 1$ (see the proof of lemma 1). Hence $aa^* = 0$ and $a = 0$. \square

Lemma 3. *Let A be a C^* -algebra of bounded linear operators on a Hilbert space H , $\{p_i\}$ be an infinite orthogonal set of projections in A with the least upper bound 1 in the algebra $B(H)$ $a \in A$. Then $a \geq 0$ if and only if for any finite subset $\{p_k\}_{k=1}^n \subset \{p_i\}$ the inequality $pap \geq 0$ holds, where $p = \sum_{k=1}^n p_k$.*

Proof. By positivity of the operator $T : a \rightarrow bab, a \in A$ for any $b \in A$, if $a \geq 0$, then for any finite subset $\{p_k\}_{k=1}^n \subset \{p_i\}$ the inequality $pap \geq 0$ holds.

Conversely, let $a \in A$. Suppose that for any finite subset $\{p_k\}_{k=1}^n \subset \{p_i\}$ the inequality $pap \geq 0$ holds, where $p = \sum_{k=1}^n p_k$.

Let $a = c + id$, for some nonzero self-adjoint elements c, d in A . Then $(p_i + p_j)(c + id)(p_i + p_j) = (p_i + p_j)c(p_i + p_j) + i(p_i + p_j)d(p_i + p_j) \geq 0$ for all i, j . In this case the elements $(p_i + p_j)c(p_i + p_j)$ and $(p_i + p_j)d(p_i + p_j)$ are self-adjoint. Then $(p_i + p_j)d(p_i + p_j) = 0$ and $p_i dp_j = 0$ for all i, j . Hence by lemma 2 we have $d = 0$. Therefore $a = c = c^* = a^*$, i.e. $a \in A_{sa}$. Hence, a is a nonzero self-adjoint element in A . Let $b_n^\alpha = \sum_{k,l=1}^n p_k^\alpha a p_l^\alpha$ for all natural numbers n and finite subsets $\{p_k^\alpha\}_{k=1}^n \subset \{p_i\}$. Then the set (b_n^α) ultraweakly converges to the element a .

Indeed, we have $A \subseteq B(H)$. Let $\{q_\xi\}$ be a maximal orthogonal set of minimal projections of the algebra $B(H)$ such, that $p_i = \sup_\eta q_\eta$, for some subset $\{q_\eta\} \subset \{q_\xi\}$, for any i . For arbitrary projections q and p in $\{q_\xi\}$ there exists a number $\lambda \in \mathbb{C}$ such, that $qap = \lambda u$, where u is an isometry in $B(H)$, satisfying the conditions $q = uu^*, p = u^*u$. Let $q_\xi q_\eta = q_\xi, q_{\xi\eta}$ be such element that $q_\xi = q_{\xi\eta} q_{\xi\eta}^*, q_\eta = q_{\xi\eta}^* q_{\xi\eta}$ for all different ξ and η . Then, let $\{\lambda_{\xi\eta}\}$ be a set of numbers such that $q_\xi a q_\eta = \lambda_{\xi\eta} q_{\xi\eta}$ for all ξ, η . In this case, since $q_\xi a a^* q_\xi = q_\xi (\sum_\eta \lambda_{\xi\eta} \bar{\lambda}_{\xi\eta}) q_\xi < \infty$, the quantity of nonzero numbers of the set $\{\lambda_{\xi\eta}\}_\eta$ (ξ -th string of the infinite dimensional matrix $\{\lambda_{\xi\eta}\}_{\xi\eta}$) is not greater then the countable cardinal number and the sequence (λ_n^ξ) of all these nonzero numbers converges to zero. Let v_{q_ξ} be a vector of the Hilbert space H , which generates the minimal projection q_ξ . Then the set $\{v_{q_\xi}\}$ forms a complete orthonormal system of the space H . Let v be an arbitrary vector of the space H and μ_ξ be a coefficient of Fourier of the vector v , corresponding to v_{q_ξ} , in relative to the complete orthonormal system $\{v_{q_\xi}\}$. Then, since $\sum_\xi \mu_\xi \bar{\mu}_\xi < \infty$, then the quantity of all nonzero elements of the set $\{\mu_\xi\}_\xi$ is not greater then the countable cardinal number and the sequence (μ_n) of all these nonzero numbers converges to zero.

Let ν_ξ be the ξ -th coefficient of Fourier (corresponding to v_{q_ξ}) of the vector $a(v) \in H$ in relative to the complete orthonormal system $\{v_{q_\xi}\}$. Then $\nu_\xi = \sum_\eta \lambda_{\xi\eta} \mu_\eta$ and the scalar product $\langle a(v), v \rangle$ is equal to the sum $\sum_\xi \nu_\xi \mu_\xi$. Since the element $a(v)$ belongs to H we have the quantity of all nonzero elements in the set $\{\nu_\xi\}_\xi$ is not greater then the countable cardinal number and the sequence (ν_n) of all these nonzero numbers converges to zero.

Let ε be an arbitrary positive number. Then, since quantity of nonzero numbers of the sets $\{\mu_\xi\}_\xi$ and $\{\nu_\xi\}_\xi$ is not greater then the countable cardinal number, and $\sum_\xi \nu_\xi \bar{\nu}_\xi < \infty, \sum_\xi \mu_\xi \bar{\mu}_\xi < \infty$, then there exists $\{f_k\}_{k=1}^l \subset \{p_i\}$ such, that for the

set of indexes $\Omega_1 = \{\xi : \exists p \in \{f_k\}_{k=1}^l, q_\xi \leq p\}$ we have

$$|\sum_{\xi} \nu_{\xi} \mu_{\xi} - \sum_{\xi \in \Omega_1} \nu_{\xi} \mu_{\xi}| < \varepsilon.$$

Then, since quantity of nonzero numbers of the sets $\{\mu_{\xi}\}_{\xi}$ and $\{\lambda_{\xi\eta}\}_{\eta}$ is not greater than the countable cardinal number, and $\sum_{\eta} \lambda_{\xi\eta} \bar{\lambda}_{\xi\eta} < \infty$, $\sum_{\xi} \mu_{\xi} \bar{\mu}_{\xi} < \infty$, then there exists $\{e_k\}_{k=1}^m \subset \{p_i\}$ such, that for the set of indexes $\Omega_2 = \{\xi : \exists p \in \{e_k\}_{k=1}^m, q_{\xi} \leq p\}$ we have

$$|\sum_{\eta} \lambda_{\xi\eta} \mu_{\eta} - \sum_{\eta \in \Omega_2} \lambda_{\xi\eta} \mu_{\eta}| < \varepsilon.$$

Hence for the finite set $\{p_k\}_{k=1}^n = \{f_k\}_{k=1}^l \cup \{e_k\}_{k=1}^m$ and the set $\Omega = \{\xi : \exists p \in \{p_k\}_{k=1}^n, q_{\xi} \leq p\}$ of indexes we have

$$|\sum_{\xi} \nu_{\xi} \mu_{\xi} - \sum_{\xi \in \Omega} (\sum_{\eta \in \Omega} \lambda_{\xi\eta} \mu_{\eta}) \mu_{\xi}| < \varepsilon.$$

At the same time, $\langle (\sum_{kl=1}^n p_k a p_l)(v), v \rangle = \sum_{\xi \in \Omega} (\sum_{\eta \in \Omega} \lambda_{\xi\eta} \mu_{\eta}) \mu_{\xi}$. Therefore,

$$|\langle a(v), v \rangle - \langle (\sum_{kl=1}^n p_k a p_l)(v), v \rangle| < \varepsilon.$$

Hence, since the vector v and the number ε are chosen arbitrarily, we have the net (b_n^{α}) ultraweakly converges to the element a .

We have there exists a maximal orthogonal set $\{e_{\xi}\}$ of minimal projections of the algebra $B(H)$ of all bounded linear operators on H , such that the element a and the set $\{e_{\xi}\}$ belong to some maximal commutative subalgebra A_o of the algebra $B(H)$. We have for any finite subset $\{p_k\}_{k=1}^n \subset \{p_i\}$ and $e \in \{e_{\xi}\}$ the inequality $e(\sum_{kl=1}^n p_k a p_l)e \geq 0$ holds by the positivity of the operator $T : b \rightarrow ebe, b \in A$.

By the previous part of the proof the net $(e_{\xi} b_n^{\alpha} e_{\xi})_{\alpha n}$ ultraweakly converges to the element $e_{\xi} a e_{\xi}$ for any index ξ . Then we have $e_{\xi} b_n^{\alpha} e_{\xi} \geq 0$ for all n and α . Therefore, the ultraweak limit $e_{\xi} a e_{\xi}$ of the net $(e_{\xi} b_n^{\alpha} e_{\xi})_{\alpha n}$ is a nonnegative element. Hence, $e_{\xi} a e_{\xi} \geq 0$. Therefore, since e_{ξ} is chosen arbitrarily then $a \geq 0$. \square

Lemma 4. *Let A be a C^* -algebra of bounded linear operators on a Hilbert space H , $\{p_i\}$ be an infinite orthogonal set of projections in A with the least upper bound 1 in the algebra $B(H)$ $a \in A$. Then*

$$\|a\| = \sup\{\|\sum_{kl=1}^n p_k a p_l\| : n \in N, \{p_k\}_{k=1}^n \subseteq \{p_i\}\}.$$

Proof. The inequality $-\|a\|1 \leq a \leq \|a\|1$ holds. Then $-\|a\|p \leq pap \leq \|a\|p$ for all natural number n and finite subset $\{p_k^{\alpha}\}_{k=1}^n \subset \{p_i\}$, where $p = \sum_{k=1}^n p_k$. Therefore

$$\|a\| \geq \sup\{\|\sum_{kl=1}^n p_k a p_l\| : n \in N, \{p_k\}_{k=1}^n \subseteq \{p_i\}\}.$$

At the same time, since the finite subset $\{p_k\}_{k=1}^n$ of $\{p_i\}$ is chosen arbitrarily and by lemma 6 we have

$$\|a\| = \sup\{\|\sum_{kl=1}^n p_k a p_l\| : n \in N, \{p_k\}_{k=1}^n \subseteq \{p_i\}\}.$$

Otherwise, if

$$\|a\| > \lambda = \sup\{\|\sum_{kl=1}^n p_k a p_l\| : n \in N, \{p_k\}_{k=1}^n \subseteq \{p_i\}\},$$

then by lemma 3 $-\lambda 1 \leq a \leq \lambda 1$. But the last inequality is a contradiction. \square

Lemma 5. *Let A be a C^* -algebra of bounded linear operators on a Hilbert space H , $\{p_i\}$ be an infinite orthogonal set of projections in A with the least upper bound 1 in the algebra $B(H)$, and let $\mathcal{A} = \{\{p_i a p_j\} : a \in A\}$. Then,*

1) *the vector space \mathcal{A} is a unit order space respect to the order $\{p_i a p_j\} \geq 0$ ($\{p_i a p_j\} \geq 0$ if for any finite subset $\{p_k\}_{k=1}^n \subset \{p_i\}$ the inequality $p a p \geq 0$ holds, where $p = \sum_{k=1}^n p_k$, and the norm*

$$\|\{p_i a p_j\}\| = \sup\{\|\sum_{kl=1}^n p_k a p_l\| : n \in N, \{p_k\}_{k=1}^n \subseteq \{p_i\}\}.$$

2) *the algebra A and the unit order space \mathcal{A} can be identified as unit order spaces in the sense of the map*

$$\mathcal{I} : a \in A \rightarrow \{p_i a p_j\} \in \mathcal{A}.$$

Proof. This lemma follows by lemmas 1, 3 and 4. \square

Remark. Observe that by lemma 4 the order and the norm in the unit order space $\mathcal{A} = \{\{p_i a p_j\} : a \in A\}$ can be defined as follows to: $\{p_i a p_j\} \geq 0$, if $a \geq 0$; $\|\{p_i a p_j\}\| = \|a\|$. By lemmas 3 and 4 they are equivalent to the order and the norm, defined in lemma 5, correspondingly.

Let A be a C^* -algebra, $\{p_i\}$ be a countable orthogonal set of equivalent projections in A such that $\sup_i p_i = 1$ and

$$\sum_{ij}^o p_i A p_j = \{\{a_{ij}\} : \text{for any indexes } i, j, a_{ij} \in p_i A p_j, \text{ and}$$

$$\|\sum_{k=1, \dots, i-1} (a_{ki} + a_{ik}) + a_{ii}\| \rightarrow 0 \text{ at } i \rightarrow \infty\}.$$

If we introduce the componentwise algebraic operations in this set, then $\sum_{ij}^o p_i A p_j$ becomes a vector space. Also, note that $\sum_{ij}^o p_i A p_j$ is a vector subspace of \mathcal{A} . Observe that $\sum_{ij}^o p_i A p_j$ is a normed subspace of the algebra \mathcal{A} and $\|\sum_{i,j=1}^n a_{ij} - \sum_{i,j=1}^{n+1} a_{ij}\| \rightarrow 0$ at $n \rightarrow \infty$ for any $\{a_{ij}\} \in \sum_{ij}^o p_i A p_j$.

Let $\sum_{ij}^o a_{ij} := \lim_{n \rightarrow \infty} \sum_{i,j=1}^n a_{ij}$, for any $\{a_{ij}\} \in \sum_{ij}^o p_i A p_j$, and

$$C^*(\{p_i A p_j\}_{ij}) := \{\sum_{ij}^o a_{ij} : \{a_{ij}\} \in \sum_{ij}^o p_i A p_j\}.$$

Then $C^*(\{p_i A p_j\}_{ij}) \subseteq A$. By lemma 5 A and \mathcal{A} can be identified. We observe that, the normed spaces $\sum_{ij}^o p_i A p_j$ and $C^*(\{p_i A p_j\}_{ij})$ can also be identified. Further, without loss of generality we will use these identifications.

Theorem 6. *Let A be a unital C^* -algebra, $\{p_i\}$ be a countable orthogonal set of equivalent projections in A and $\sup_i p_i = 1$. Then $\sum_{ij}^o p_i A p_j$ is a C^* -subalgebra of A with the componentwise algebraic operations, the associative multiplication and the norm.*

Proof. We have $\sum_{ij}^o p_i A p_j$ is a normed subspace of the algebra A .

Let (a_n) be a sequence of elements in $\sum_{ij}^o p_i A p_j$ such that (a_n) norm converges to some element $a \in A$. We have $p_i a_n p_j \rightarrow p_i a p_j$ at $n \rightarrow \infty$ for all i and j . Hence $p_i a p_j \in p_i A p_j$ for all i, j . Let $b^n = \sum_{k=1}^n (p_{n-1} a p_k + p_k a p_{n-1}) + p_n a p_n$ and $c_m^n = \sum_{k=1}^n (p_{n-1} a_m p_k + p_k a_m p_{n-1}) + p_n a_m p_n$, for any n . Then $c_m^n \rightarrow b^n$ at $m \rightarrow \infty$. It should be proven that $\|b_n\| \rightarrow 0$ at $n \rightarrow \infty$.

Let $\varepsilon \in \mathbb{R}_+$. Then there exists m_o such that $\|a - a_m\| < \varepsilon$ for any $m > m_o$. Also for all n and $\{p_k\}_{k=1}^n \subset \{p_i\}$ $\|(\sum_{k=1}^n p_k)(a - a_m)(\sum_{k=1}^n p_k)\| < \varepsilon$. Hence $\|b^n - c_m^n\| < 2\varepsilon$ for any $m > m_o$. At the same time, $\|b^n - c_{m_1}^n\| < 2\varepsilon$, $\|b^n - c_{m_2}^n\| < 2\varepsilon$ for all $m_o < m_1, m_2$. Since $(a_n) \subset \sum_{ij}^o p_i A p_j$ then for any m $\|c_m^n\| \rightarrow 0$ at $n \rightarrow \infty$. Hence, since $\|c_{m_1}^n\| \rightarrow 0$ and $\|c_{m_2}^n\| \rightarrow 0$ at $n \rightarrow \infty$, then there exists n_o such that $\|c_{m_1}^n\| < \varepsilon$, $\|c_{m_2}^n\| < \varepsilon$ and $\|c_{m_1}^n + c_{m_2}^n\| < 2\varepsilon$ for any $n > n_o$. Then $\|2b_n\| = \|b^n - c_{m_1}^n + c_{m_1}^n + c_{m_2}^n + b^n - c_{m_2}^n\| \leq \|b^n - c_{m_1}^n\| + \|c_{m_1}^n + c_{m_2}^n\| + \|b^n - c_{m_2}^n\| < 2\varepsilon + 2\varepsilon + 2\varepsilon = 6\varepsilon$ for any $n > n_o$, i.e. $\|b_n\| < 3\varepsilon$ for any $n > n_o$. Since ε is arbitrarily chosen then $\|b_n\| \rightarrow 0$ at $n \rightarrow \infty$. Therefore $a \in \sum_{ij}^o p_i A p_j$. Since the sequence (a_n) is arbitrarily chosen then $\sum_{ij}^o p_i A p_j$ is a Banach space.

Let $\{a_{ij}\}, \{b_{ij}\}$ be arbitrary elements of the Banach space $\sum_{ij}^o p_i A p_j$. Let $a_m = \sum_{kl=1}^m a_{kl}$, $b_m = \sum_{kl=1}^m b_{kl}$ for all natural numbers m . We have the sequence (a_m) converges to $\{a_{ij}\}$ and the sequence (b_m) converges to $\{b_{ij}\}$ in $\sum_{ij}^o p_i A p_j$. Also for all n and m $a_m b_n \in \sum_{ij}^o p_i A p_j$. Then for any n the sequence $(a_m b_n)$ converges to $\{a_{ij}\} b_n$ at $m \rightarrow \infty$. Hence $\{a_{ij}\} b_n \in \sum_{ij}^o p_i A p_j$. Note that $\sum_{ij}^o p_i A p_j \subseteq A$. Therefore for any $\varepsilon \in \mathbb{R}_+$ there exists n_o such that $\|\{a_{ij}\} b_{n+1} - \{a_{ij}\} b_n\| \leq \|\{a_{ij}\}\| \|b_{n+1} - b_n\| \leq \varepsilon$ for any $n > n_o$. Hence the sequence $(\{a_{ij}\} b_n)$ converges to $\{a_{ij}\} \{b_{ij}\}$ at $n \rightarrow \infty$. Since $\sum_{ij}^o p_i A p_j$ is a Banach space then $\{a_{ij}\} \{b_{ij}\} \in \sum_{ij}^o p_i A p_j$. Since $\sum_{ij}^o p_i A p_j \subseteq A$, then $\sum_{ij}^o p_i A p_j$ is a C^* -algebra. \square

Let H be an infinite dimensional Hilbert space, $B(H)$ be the algebra of all bounded linear operators. Let $\{p_i\}$ be a countable orthogonal set of equivalent projections in $B(H)$ and $\sup_i p_i = 1$. Let $\{\{p_j^\xi\}_j\}_i$ be the set of infinite subsets of $\{p_i\}$ such that for all distinct ξ and η $\{p_j^\xi\}_j \cap \{p_j^\eta\}_j = \emptyset$, $|\{p_j^\xi\}_j| = |\{p_j^\eta\}_j|$ and $\{p_i\} = \cup_i \{p_j^\xi\}_j$. Then let $q_i = \sup_j p_j^\xi$ in $B(H)$, for all i . Then $\sup_i q_i = 1$ and $\{q_i\}$ be a countable orthogonal set of equivalent projections. Then we say that the countable orthogonal set $\{q_i\}$ of equivalent projections is defined by the set $\{p_i\}$ in $B(H)$. We have the next corollary.

Corollary 7. *Let A be a unital C^* -algebra of bounded linear operators in a Hilbert space H , $\{p_i\}$ be a countable orthogonal set of equivalent projections in A and $\sup_i p_i = 1$. Let $\{q_i\}$ be a countable orthogonal set of equivalent projections in $B(H)$ defined by the set $\{p_i\}$ in $B(H)$. Then $\sum_{ij}^o q_i A q_j$ is a C^* -subalgebra of the algebra A .*

Proof. Let $\{\{p_j^\xi\}_j\}_i$ be the set of infinite subsets of $\{p_i\}$ such that for all distinct ξ and η $\{p_j^\xi\}_j \cap \{p_j^\eta\}_j = \emptyset$, $|\{p_j^\xi\}_j| = |\{p_j^\eta\}_j|$ and $\{p_i\} = \cup_i \{p_j^\xi\}_j$. Then let $q_i = \sup_j p_j^\xi$ in $B(H)$, for all i . Then we have for all i and j $q_i A q_j = \{\{p_\xi^i a p_\eta^j\}_{\xi\eta} : a \in A\}$. Hence $q_i A q_j \subset A$ for all i and j .

The rest part of the proof is the repeating of the proof of theorem 6. \square

Example. 1. Let \mathcal{M} be the closure on the norm of the inductive limit \mathcal{M}_o of the inductive system

$$C \rightarrow M_2(C) \rightarrow M_3(C) \rightarrow M_4(C) \rightarrow \dots,$$

where $M_n(C)$ is mapped into the upper left corner of $M_{n+1}(C)$. Then \mathcal{M} is a C^* -algebra ([1]). The algebra \mathcal{M} contains the minimal projections of the form e_{ii} , where e_{ij} is an infinite dimensional matrix, whose (i, i) -th component is 1 and the rest components are zeros. These projections form the countable orthogonal set $\{e_{ii}\}_{i=1}^\infty$ of minimal projections. Let

$$M_n^o(\mathbb{C}) = \left\{ \sum_{ij} \lambda_{ij} e_{ij} : \text{for any indexes } i, j, \lambda_{ij} \in \mathbb{C}, \text{ and} \right.$$

$$\left. \left\| \sum_{k=1, \dots, i-1} (\lambda_{ki} e_{ki} + \lambda_{ik} e_{ik}) + \lambda_{ii} e_{ii} \right\| \rightarrow 0 \text{ at } i \rightarrow \infty \right\}.$$

Then $\mathbb{C} \cdot 1 + M_n^o(\mathbb{C}) = \mathcal{M}$ (see [2]) and by theorem 6 $M_n^o(\mathbb{C})$ is a simple C^* -algebra. Note that there exists a mistake in the formulation of theorem 3 in [2]. $\mathbb{C} \cdot 1 + M_n^o(\mathbb{C})$ is a C^* -algebra. But the algebra $\mathbb{C} \cdot 1 + M_n^o(\mathbb{C})$ is not simple. Because $\mathbb{C} \cdot 1 + M_n^o(\mathbb{C}) \neq M_n^o(\mathbb{C})$ and $M_n^o(\mathbb{C})$ is an ideal of the algebra $\mathbb{C} \cdot 1 + M_n^o(\mathbb{C})$, i.e. $[\mathbb{C} \cdot 1 + M_n^o(\mathbb{C})] \cdot M_n^o(\mathbb{C}) \subseteq M_n^o(\mathbb{C})$.

2. There exist a C^* -algebra A and different countable orthogonal sets $\{e_i\}$ and $\{f_i\}$ of equivalent projections in A such that $\sup_i e_i = 1$, $\sup_i f_i = 1$, $\sum_{ij}^o e_i A e_j \neq \sum_{ij}^o f_i A f_j$. Indeed, let H be an infinite dimensional Hilbert space, $B(H)$ be the algebra of all bounded linear operators. Let $\{p_i\}$ be a countable orthogonal set of equivalent projections in $B(H)$ and $\sup_i p_i = 1$. Then $\sum_{ij}^o p_i B(H) p_j \subset B(H)$. Let $\{\{p_j^i\}_j\}_i$ be the set of infinite subsets of $\{p_i\}$ such that for all distinct ξ and η $\{p_j^\xi\}_j \cap \{p_j^\eta\}_j = \emptyset$, $|\{p_j^\xi\}_j| = |\{p_j^\eta\}_j|$ and $\{p_i\} = \cup_i \{p_j^i\}_j$. Then let $q_i = \sup_j p_j^i$ for all i . Then $\sup_i q_i = 1$ and $\{q_i\}$ be a countable orthogonal set of equivalent projections. We assert that $\sum_{ij}^o p_i B(H) p_j \neq \sum_{ij}^o q_i B(H) q_j$. Indeed, let $\{x_{ij}\}$ be a set of matrix units constructed by the infinite set $\{p_j^1\}_j \in \{\{p_j^i\}_j\}_i$, i.e. for all i, j , $x_{ij} x_{ij}^* = p_i^1$, $x_{ij}^* x_{ij} = p_j^1$, $x_{ii} = p_i^1$. Then the von Neumann algebra \mathcal{N} generated by the set $\{x_{ij}\}$ is isometrically isomorphic to $B(\mathcal{H})$ for some Hilbert space \mathcal{H} . We note that \mathcal{N} is not subset of $\sum_{ij}^o p_i B(H) p_j$. At the same time, $\mathcal{N} \subseteq \sum_{ij}^o q_i B(H) q_j$ and $\sum_{ij}^o p_i^1 \mathcal{N} p_j^1 \subseteq \sum_{ij}^o p_i B(H) p_j$.

Theorem 8. Let A be a unital simple C^* -algebra of bounded linear operators in a Hilbert space H , $\{p_i\}$ be a countable orthogonal set of equivalent projections in A and $\sup_i p_i = 1$. Let $\{q_i\}$ be a countable orthogonal set of equivalent projections in $B(H)$ defined by the set $\{p_i\}$ in $B(H)$. Then $\sum_{ij}^o q_i A q_j$ is a simple C^* -algebra.

Proof. By theorem 6 $\sum_{ij}^o p_i A p_j$ is a C^* -algebra. Let $\{\{p_j^i\}_j\}_i$ be the set of infinite subsets of $\{p_i\}$ such that for all distinct ξ and η $\{p_j^\xi\}_j \cap \{p_j^\eta\}_j = \emptyset$, $|\{p_j^\xi\}_j| = |\{p_j^\eta\}_j|$ and $\{p_i\} = \cup_i \{p_j^i\}_j$. Then let $q_i = \sup_j p_j^i$ in $B(H)$, for all i . Then we have for all i and j $q_i A q_j = \{\{p_\xi^i a p_\xi^j\} : a \in A\}$. Hence $q_i A q_j \subset A$ for all i and j . By corollary 7 $\sum_{ij}^o q_i A q_j$ is a C^* -algebra.

Since projections of the set $\{p_i\}$ pairwise equivalent then the projection q_i is equivalent to $1 \in A$ for any i . Hence $q_i A q_i \cong A$ and $q_i A q_i$ is a simple C^* -algebra for any i .

Let q be an arbitrary projection in $\{q_i\}$. Then qAq is a subalgebra of $\sum_{ij}^o q_i A q_j$. Let I be a closed ideal of the algebra $\sum_{ij}^o q_i A q_j$. Then $IqAq \subset I$ and $Iq \cdot qAq \subset Iq$. Therefore $qIqqAq \subseteq qIq$, that is qIq is a closed ideal of the subalgebra qAq . Since qAq is simple then $qIq = qAq$.

Let q_1, q_2 be arbitrary projections in $\{q_i\}$. We assert that $q_1 I q_2 = q_1 A q_2$ and $q_2 I q_1 = q_2 A q_1$. Indeed, we have the projection $q_1 + q_2$ is equivalent to $1 \in A$. Let $e = q_1 + q_2$. Then $eAe \cong A$ and eAe is a simple C^* -algebra. At the same time we have eAe is a subalgebra of $\sum_{ij}^o q_i A q_j$ and I is an ideal of $\sum_{ij}^o q_i A q_j$. Hence $IeAe \subset I$ and $Ie \cdot eAe \subset Ie$. Therefore $eIeeAe \subseteq eIe$, that is eIe is a closed ideal of the subalgebra eAe . Since eAe is simple then $eIe = eAe$. Hence $q_1 I q_2 = q_1 A q_2$ and $q_2 I q_1 = q_2 A q_1$. Therefore $q_i I q_j = q_i A q_j$ for all i and j . We have I is norm closed. Hence $I = \sum_{ij}^o q_i A q_j$, i.e. $\sum_{ij}^o q_i A q_j$ is a simple C^* -algebra. \square

2. APPLICATIONS

Theorem 9. *Let \mathcal{N} be a W^* -factor of type II_∞ of bounded linear operators in a Hilbert space H , $\{p_i\}$ be a countable orthogonal set of equivalent projections in \mathcal{N} and $\sup_i p_i = 1$. Then for any countable orthogonal set $\{q_i\}$ of equivalent projections in $B(H)$ defined by the set $\{p_i\}$ in $B(H)$ the C^* -algebra $\sum_{ij}^o q_i \mathcal{N} q_j$ is a C^* -factor with a nonzero finite and an infinite projection. In this case $\sum_{ij}^o q_i \mathcal{N} q_j$ is not a von Neumann algebra.*

Proof. By the definition of the set $\{q_i\}$ we have $\sup_i q_i = 1$ and $\{q_i\}$ be a countable orthogonal set of equivalent *infinite* projections. By theorem 6 we have $\sum_{ij}^o q_i \mathcal{N} p_j$ is a C^* -subalgebra of \mathcal{N} . Let q be a nonzero finite projection of \mathcal{N} . Then there exists a projection $p \in \{p_i\}$ such that $qp \neq 0$. We have $q\mathcal{N}q$ is a finite von Neumann algebra. Let $x = pq$. Then $x\mathcal{N}x^*$ is a weakly closed C^* -subalgebra. Note that the algebra $x\mathcal{N}x^*$ has a center-valued faithful trace. Let e be a nonzero projection of the algebra $x\mathcal{N}x^*$. Then $ep = e$ and $e \in p\mathcal{N}p$. Hence $e \in \sum_{ij}^o q_i \mathcal{N} q_j$. We have the weak closure of $\sum_{ij}^o q_i \mathcal{N} q_j$ in the algebra \mathcal{N} coincides with this algebra \mathcal{N} . Then by the weak continuity of the multiplication $\sum_{ij}^o q_i \mathcal{N} q_j$ is a factor. Note since $1 \notin \sum_{ij}^o q_i \mathcal{N} q_j$ then $\sum_{ij}^o q_i \mathcal{N} q_j$ is not weakly closed in \mathcal{N} . Hence the C^* -factor $\sum_{ij}^o q_i \mathcal{N} q_j$ is not a von Neumann algebra. \square

Remark. Note that, in the article [3] a simple C^* -algebra with an infinite and a nonzero finite projection have been constructed by M.Rørdam. In the next corollary we construct a simple purely infinite C^* -algebra. Note that simple purely infinite C^* -algebras are considered and investigated, in particular, in [4] and [5].

Theorem 10. *Let \mathcal{N} be a W^* -factor of type III of bounded linear operators in a Hilbert space H . Then for any countable orthogonal set $\{p_i\}$ of equivalent projections in \mathcal{N} such that $\sup_i p_i = 1$, $\sum_{ij}^o p_i \mathcal{N} p_j$ is a simple purely infinite C^* -algebra. In this case $\sum_{ij}^o p_i \mathcal{N} p_j$ is not a von Neumann algebra.*

Proof. Let p_{i_0} be a projection in $\{p_i\}$. We have the projection p_{i_0} can be exhibited as a least upper bound of a countable orthogonal set $\{p_{i_0}^j\}_j$ of equivalent projections in \mathcal{N} . Then for any i the projection p_i has a countable orthogonal set $\{p_i^j\}_j$ of equivalent projections in \mathcal{N} such that the set $\bigcup_i \{p_i^j\}_j$ is a countable orthogonal set of equivalent projections in \mathcal{N} . Hence the set $\{p_i\}$ is defined by the set $\bigcup_i \{p_i^j\}_j$ in $B(H)$ (in \mathcal{N}). Hence by theorem 8 $\sum_{ij}^o p_i \mathcal{N} p_j$ is a simple C^* -algebra. Note since

$1 \notin \sum_{ij}^o p_i \mathcal{N} p_j$ then $\sum_{ij}^o p_i \mathcal{N} p_j$ is not weakly closed in \mathcal{N} . Hence $\sum_{ij}^o p_i \mathcal{N} p_j$ is not a von Neumann algebra.

Suppose there exists a nonzero finite projection q in $\sum_{ij}^o p_i \mathcal{N} p_j$. Then there exists a projection $p \in \{p_i\}$ such that $qp \neq 0$. We have $q(\sum_{ij}^o p_i \mathcal{N} p_j)q$ is a finite C^* -algebra. Let $x = pq$. Then $x\mathcal{N}x^*$ is a C^* -subalgebra. Moreover $x\mathcal{N}x^*$ is weakly closed and $x\mathcal{N}x^* \subset p\mathcal{N}p$. Hence $x\mathcal{N}x^*$ has a center-valued faithful trace. Then $x\mathcal{N}x^*$ is a finite von Neumann algebra with a center-valued faithful normal trace. Let e be a nonzero projection of the algebra $x\mathcal{N}x^*$. Then $ep = e$ and $e \in p\mathcal{N}p$. Hence $e \in \mathcal{N}$. This is a contradiction. \square

Example. Let H be a separable Hilbert space and $B(H)$ the algebra of all bounded linear operators on H . Let $\{q_i\}$ be a maximal orthogonal set of equivalent minimal projections in $B(H)$. Then $\sum_{ij}^o q_i B(H) q_j$ is a two sided closed ideal of the algebra $B(H)$. Using the set $\{q_i\}$ we construct a countable orthogonal set $\{p_i\}$ of equivalent infinite projections such that $\sup_i p_i = 1$. Let $\{\{q_j^{i_1}\}_j\}_i$ be the countable set of countable subsets of $\{q_i\}$ such that for all distinct i_1 and i_2 $\{q_j^{i_1}\}_j \cap \{q_j^{i_2}\}_j = \emptyset$ and $\{q_i\} = \cup_i \{q_j^{i_1}\}_j$. Then let $p_i = \sup_j q_j^{i_1}$ for all i . Then $\sup_i p_i = 1$ and $\{p_i\}$ is a countable orthogonal set of equivalent infinite projections in $B(H)$ defined by $\{q_i\}$ in $B(H)$.

Let $\{q_{nm}^{ij}\}$ be the set of matrix units constructed by the set $\{\{q_j^{i_1}\}_j\}_i$, i.e. $q_{nm}^{ij} q_{nm}^{ij*} = q_n^i$, $q_{nm}^{ij*} q_{nm}^{ij} = q_m^j$, $q_{nn}^{ii} = q_n^i$ for all i, j, n, m . Then let $a = \{a_{nm}^{ij} q_{nm}^{ij}\}$ be the decomposition of the element $a \in B(H)$, where the components a_{nm}^{ij} are defined as follows

$$a_{11}^{11} = \lambda, a_{12}^{21} = \lambda, a_{13}^{31} = \lambda, \dots, a_{1n}^{n1} = \lambda, \dots,$$

and the rest components a_{nm}^{ij} are zero, i.e. $a_{nm}^{ij} = 0$. Then $p_1 a = a$. Then since $a \notin \sum_{ij}^o p_i B(H) p_j$ and $p_1 \in \sum_{ij}^o p_i B(H) p_j$ then $\sum_{ij}^o p_i B(H) p_j$ is not an ideal of $B(H)$. But by theorem 6 $\sum_{ij}^o p_i B(H) p_j$ is a C^* -algebra. Hence there exists a C^* -algebra A with an orthogonal set $\{p_i\}$ of equivalent projections such that $\sum_{ij}^o p_i A p_j$ is not an ideal of A .

REFERENCES

- [1] O.Bratteli. *Inductive limits of finite dimensional C^* -algebras*. Trans. Amer. Math. Soc. 171, pp. 195-234. (1972)
- [2] F. N. Arzikulov. *Infinite order and norm decompositions of C^* -algebras*. Int. Journal of Math. Analysis, Vol. 2, no. 5-8, pp. 255-262.(2008) (www.m-hikari.com)
- [3] M.Rørdam. *A simple C^* -algebra with a finite and an infinite projection*. Acta Math. 191. p. 109-142. (2003)
- [4] N.C.Philips. *A Classification Theorem for Nuclear Purely Infinite Simple C^* -Algebras*. Documenta Math. No. 5. p. 49-114. (2000)
- [5] M.Rørdam. *Classification of Nuclear C^* -algebras*. Springer, Encyclopaedia of Mathematical Sciences, Vol. 126, (2001).

E-mail address: arzikulovfn@yahoo.com

Current address: Institute of Mathematics and Information technologies, Tashkent